

World Maritime University The Maritime Commons: Digital Repository of the World Maritime University

NSBWO

Project Reports

8-12-2013

Biofouling : a means of aquatic species transfer

Anne Bouyssou

World Maritime University

Josefin Madjidian

World Maritime University, jam@wmu.se

Follow this and additional works at: <http://commons.wmu.se/nsbwo>



Part of the [Ecology and Evolutionary Biology Commons](#), [Environmental Health and Protection Commons](#), [Marine Biology Commons](#), and the [Transportation Commons](#)

Recommended Citation

Bouyssou, Anne and Madjidian, Josefin, "Biofouling : a means of aquatic species transfer" (2013). NSBWO. 3.
<http://commons.wmu.se/nsbwo/3>

This Report Open Access is brought to you courtesy of Maritime Commons. Open Access items may be downloaded for non-commercial, fair use academic purposes. No items may be hosted on another server or web site without express written permission from the World Maritime University. For more information, please contact library@wmu.se.

BIOFOULING : A MEANS OF AQUATIC SPECIES TRANSFER

Anne Bouyssou & Josefin Madjidian
World Maritime University
PO Box 500
S-201 24 Malmö
Sweden

Reporting Period 9, Mar to Aug 2013
12-08-2013

This report should be quoted as follows

Bouyssou A and Madjidian, J.A. 2014. Biofouling: a means of aquatic species transfer. Final report. Prepared for Interreg IVB North Sea Ballast Water Opportunity project: 18pp.

Disclaimer

The contents and views contained in this report are those of the authors, and do not necessarily represent those of the Interreg IVB North Sea Region Programme. Although careful consideration is given to its content, we cannot be held responsible for its use.

TABLE OF CONTENTS

1	INTRODUCTION	3
2	WHAT IS BIOFOULING?	3
3	BIOFOULING OF SHIPS AND POTENTIAL IMPACTS	6
4	HOW TO REDUCE BIOFOULING OF SHIPS?.....	7
5	POSSIBLE SIDE-EFFECTS OF SHIPS' BIOFOULING CONTROL METHODS.....	9
6	INTERNATIONAL BIOFOULING MANAGEMENT MEASURES.....	9
7	EXAMPLES OF SPECIES INTRODUCED INTO THE NORTH SEA THROUGH SHIPS' BIOFOULING	10
8	CONCLUSION	12
9	REFERENCES	14

1 INTRODUCTION

Ships carry seawater in their ballast tanks when they are not fully loaded with cargo, in order to maintain adequate trim, draught and stability, adjust list and limit stresses on the hull. It is now well documented that the water pumped into the ship contains aquatic organisms – which can also sink to the sediments at the bottom of tanks – and that these organisms are thereby transferred from the port of origin to the destination. But it is seldom mentioned that aquatic organisms are also found on the outside of ships, attached on their hulls and appendages, as a result of **a very dynamic process called 'biofouling'**.


When comparing the two main introduction pathways associated with ships, namely ballast water and biofouling, some scientists (Drake & Lodge, 2007) regard biofouling as presenting **a higher risk of species introduction than ballast water**. However, predominance of invasive species vectors appears to differ for different regions (Gollasch, *et al.*, 2010).

2 WHAT IS BIOFOULING?

Biofouling is defined as the “accumulation of aquatic organisms such as micro-organisms, plants, and animals on surfaces and structures immersed in or exposed to the aquatic environment. [It] can include microfouling and macrofouling” (IMO, 2011). Indeed, when a clean surface is immersed in natural seawater, it immediately starts to adsorb a molecular ‘conditioning’ film primarily consisting of dissolved organic material (Jain & Bhosle, 2009).

There are more than 4,000 marine fouling species (Arai, 2009). For some, the settlement on a hard surface represents a transitory phase in their life-cycle, whereby they shift from larval stage to adult life (Callow & Callow, 2002).

The below Table provides some examples of micro- and macrofouling organisms (Table 1).



Picture source: <http://www.seos-project.eu>

Table 1: Some examples of microfouling and macrofouling organisms
(Sources: Callow & Callow, 2002; Railkin, 2004; Ministry of Agriculture and Forestry Biosecurity New Zealand, 2010a; Ministry of Agriculture and Forestry Biosecurity New Zealand, 2010b; Sorte, Williams & Zerebecki, 2010)

		TYPES	EXAMPLES
MICROFOULING ORGANISMS		Sessile bacteria	<i>Micrococcus, Pseudomonas</i>
		Diatoms	<i>Amphora</i> spp., <i>Navicula</i> sp., <i>Nitschia</i> spp.
		Micro-fungi	
		Heterotrophic flagellates	<i>Monosiga, Pteridomonas</i>
		Sarcodines	
		Sessile ciliates	
MACROFOULING ORGANISMS	Hard fouling	Barnacles	<i>Amphibalanus amphitrite, Amphibalanus reticulatus, Balanus amphitrite</i>
		Bivalves	<i>Crassostrea gigas, Mytilus</i> spp., <i>Perna canaliculus, Perna perna</i>
		Calcareous tube worms	<i>Hydroides albiceps, Hydroides elegans</i>
	Soft fouling	Algae	<i>Laminaria</i> spp. (brown alga) <i>Enteromorpha</i> spp., <i>Ulva</i> spp. (green algae) <i>Ahnfeltia</i> spp. (red alga)
		Anemones	<i>Haliplanella</i> sp.
		Ascidians	<i>Didemnum vexillum</i>
		Bryozoans	<i>Bugula neritina, Cryptosula pallasiana, Watersipora subtorquata, Zoobotryon pellucidum</i>
		Corals	
		Hydroids	<i>Obelia</i> sp.
		Sea cucumbers	
		Sponges	<i>Acanthella cavernosa</i>

It should be noted that biofouling is not purely made up of sessile organisms. It may include **mobile organisms** as well, such as crustaceans and fish (Gollasch, 2002; Ministry of Agriculture and Forestry Biosecurity New Zealand, 2010b).



Figure 1: Biofouling is an assemblage of both sessile and mobile species
(Source: Ministry of Agriculture and Forestry Biosecurity New Zealand, 2010b).

Marine organisms congregate on surfaces by a five-stage process (Callow, 2000; Zinn, Zimmerman & White, 2000; Callow & Callow, 2002; Quiniou & Compère, 2009):

- **Attachment of organic and nitrogen-based compounds, as well as salts and silica, on the surface.**

This phase lasts a few minutes. The result is that the surface becomes organically enriched with chemical compounds which occur naturally in seawater.

- **Attachment of primary colonizers, such as bacteria, algal cells and spores.**

This phase takes place within a few hours. Organisms are attracted to the surface because it offers them a source of food.

- **Excretion of extracellular polymeric substances (EPSs) by the primary colonizers.**

EPSs consist of lipids, proteins, nucleic acids, and polysaccharides. They play an important role in biofilm formation because they constitute the substratum in which micro-organisms will be encased (Flemming, *et al.*, 2000; Ahimou, *et al.*, 2007).

- **Development of a biofilm.**

The biofilm matrix is generated within a few days up to a month through cell divisions. It is composed of myriads of bacteria, protozoa, larvae, algal cells such as diatoms, and spores, separated by interstitial voids filled with water (Lewandowski, 2000). This structure, which can be up to 500 µm thick, is also known as 'microfouling' or 'slime' (Callow & Callow, 2002).

Both the internal cohesion and the adhesion of the biofilm to the colonized surface depend upon the nature and the amount of EPSs (Ahimou, *et al.*, 2007). It is noteworthy that, **when structured in a biofilm, cells are less susceptible to biocides** than in their planktonic form (Allison, *et al.*, 2000; Flemming, 2002; Russell, 2003). The reason is that the EPS matrix acts as a physical barrier that

protects biofouling communities (Donlan, 2000). Furthermore, the biofilm provides a substratum for macroorganism attachment.

- **Accumulation of macro-organisms onto the biofilm.**

Macrofouling organisms are comprised of 'hard fouling' organisms and 'soft fouling' organisms – see Table 1 (Callow & Callow, 2002).

Some fouling organisms such as barnacles, mussels and tube worms are generally found on the lower parts of the ship's underwater body, while algae commonly occupy the higher parts (Ministry of Agriculture and Forestry Biosecurity New Zealand, 2010b). Algal growth depends on light quantity and availability of space (Holmström & Kjelleberg, 2000).

Several factors, both **operational** and **environmental**, influence the above-detailed process:

- Type, colour, age and state of the antifouling coating, ship speed, trading area (Swain, *et al.*, 2006; Drake & Lodge, 2007; Schultz, *et al.*, 2011).
- Type of biofilm matrix, competition and predation among fouling communities (Callow & Callow, 2002), temperature, pH and nutrient transfer within the biofouling layers.
- Abiotic environmental aspects as temperature, solar radiation and salinity further have large impacts on species survival (e.g. Kim & Micheli, 2013; Verween, *et al.*, 2007), this specially regards ships travelling between tropic areas and colder areas or between seaports and freshwater ports.

In the early stages, organisms are easy to remove. As time goes by, they stick fast and accumulate. On some ships and in certain areas of the hull, it has been found that biofouling could reach a thickness of 30 cm (Gollasch, 2002).

3 BIOFOULING OF SHIPS AND POTENTIAL IMPACTS

Marine organisms congregate not only on the outside of ships and appendages – e.g. hulls, sea-chests, propellers, bow and stern thrusters, but also inside ships – such as on filters, heat exchangers, seawater cooling pipes, pumps and valves.

Sea-chests have lately been suggested to be hotspots for biofouling (Coutts and Taylor, 2004; Coutts and Dodgshun, 2007; Sylvester and MacIsaac, 2010, Frey *et al.*, 2014). These are protected, cavity-like structures, built into the hull of a vessel and typically covered with metal grates (Coutts *et al.*, 2003) and are typically characterized by relatively low water flows compared to higher velocities and shear stresses experienced on the exposed, flat surfaces of the hull. As a result sea-chests provide a relatively protected refuge for many fouling organisms, leading to increased survivorship and thriving communities (Coutts and Dodgshun, 2007).

It is noteworthy that all vessel categories are affected by biofouling, i.e. warships, merchant ships, fishing vessels, barges, mobile offshore units, recreational boats, etc.

Biofouling of ships is a concern for safety, economic, and environmental reasons.

Safety impacts

Biofouling may compromise safety because it hampers the proper functioning of pipework and associated appliances – for example, cooling systems, as well as the operation of navigation instruments (Callow, 2000).

Marine biofouling also accelerates biocorrosion (Meesters, *et al.*, 2003), by maintaining a continuous metal/organisms interaction, which may result in the deterioration of some ships' structures.

Economic impacts

Biofouling reduces heat transfer, makes the ship heavier, and induces greater frictional resistance on the hull and the propeller, thereby reducing fuel efficiency (Schultz, *et al.*, 2011). As an example, an increase of 100 µm in the average hull roughness augments fuel consumption approximately by 6 % (Arai, 2009).

To prevent the attachment of marine organisms, antifouling coatings have to be applied over the outside of ships. This includes removing the old coating, then repainting the hull and appendages.

Further maintenance operations might have to be carried out also inside ships – through the use of chlorine, for example (Meesters, *et al.*, 2003) – when the water flow is reduced in pipes.

In-water inspections and cleaning, as well as drydocking, generate additional costs (Callow & Callow, 2002). Removal of biofouling on ships can be executed either at predetermined intervals or when the degree of fouling makes it necessary – such as in the U.S. Navy ships (Schultz, *et al.*, 2011). Cleaning techniques involve water jets, steam, robots with rotating brushes, ultrasound or acid and base baths. It should be noted that not all methods are applicable to large vessels and mobile offshore drilling units (Zinn, *et al.*, 2000).

Environmental impacts

Biofouling might present an equivalent or even greater risk of species transfer than ballast water (Drake & Lodge, 2007), however, other studies found that ballast water is the dominating vector. Despite the application of antifouling coatings, organisms are still found on ships. They accumulate in niche areas, as well as on the surfaces where the coating is damaged, worn or inadequately applied.

As it generates frictional resistance when the vessel moves through the water, biofouling increases bunker fuel consumption, and in turn, CO₂ emissions (Zinn, *et al.*, 2000; Callow & Callow, 2002; Drake & Lodge, 2007; IMO, 2010).

4 HOW TO REDUCE BIOFOULING OF SHIPS?

Antifouling coatings

Historically, antifoulants included biocides such as lead, arsenic mercury and their organic derivatives. However, these were banned due to the environmental risks that they posed. Also the revolutionary self-polishing copolymer technique that employed a similar heavy metal toxic action to deter marine organisms, the antifoulant tributyltin (TBT), has been banned. Indeed, the use of **organotins** in antifouling paints was prohibited in 2008, with the entry into force of the International Convention on the Control of Harmful Antifouling Systems on Ships, 2001 (AFS Convention). (See chapter 6 for international biofouling management measures.)

Since 2008, two broad categories of antifouling coatings are applied on ships: those which contain toxic agents, and those that do not. The former category makes use of biocides, such as **copper** associated with booster biocides. The working principle of such paint systems is based on slow release of toxins in time (self-polishing coatings). Copper has been used for a long time as an antifouling agent (Callow & Callow, 2002). The latter, is based on biocide-free products such as silicone-based antifouling paints. A diversifying market for antifouling systems is to be expected as there is no single best solution for all ship types.




Figure 2: Two broad categories of antifouling coatings are applied on ships.

Non-toxic antifouling agents

Non-toxic paints may have a shorter lifespan than organotin paints and, consequently, may need to be applied more often (Champ, 2001; Readman, *et al.*, 2002; Chambers, *et al.*, 2006).

They might also be less efficient in preventing biofouling (Mineur, *et al.*, 2007). Another drawback of biocide-free coatings is that they may not be sufficiently robust for some deep-sea ships (Callow, 2000).

Non-toxic, fouling-resistant coatings are based on polymers designed to minimize molecular adhesive forces between the adhesives used by marine organisms and the coating. This is made through manipulation of the physicochemical and/or materials properties of the coating. In order to be able to select a surface it is necessary to understand the interfacial interactions at the molecular level. As a result, a great challenge is the vast diversity of fouling organisms and the range of their adhesion mechanisms (including adhesives) (Callow & Callow, 2011).

Natural antifouling surfaces further inspire researchers whilst searching for new coating designs. For example, marine invertebrates such as sponges and corals usually remain remarkably free from settlement by fouling organisms. Indeed, sponge-derived anti-fouling molecules have been found to inhibit the settlement of barnacle larvae (Hellio, *et al.*, 2005), inhibit fouling by macroalgae (Kubaneck, *et al.*, 2002), or repel the blue mussel *Mytilus edulis galloprovincialis* (Sera, *et al.*, 1999). There is much research to be done within this field.

Hull maintenance

Both biocidal and biocide-free antifouling coatings may contain harmful substances that pose a contamination risk if released into the environment. Therefore, application, maintenance and removal of anti-fouling coatings on vessels in maintenance facilities/dry-docking or in-water can result in contamination of the aquatic environment. For example, toxic paint particles can be released into the marine environment if ship hull cleaning procedures are improperly regulated and contained, and the release of biofouling organisms during cleaning can facilitate the spread of invasive aquatic species. Decisions on the appropriate management option for a specific ship will be influenced by many factors, including the species present, the level of fouling, and the time a vessel spends in a recipient region (Hopkins & Forrest, 2008).

5 POSSIBLE SIDE-EFFECTS OF SHIPS' BIOFOULING CONTROL METHODS

Leaching of biocides contained in antifouling coatings

The use of **organotin compounds**, such as tributyltin (TBT), in antifouling paints has produced damaging effects on marine life. In the 1980s, it was discovered that TBT can cause gastropods and bivalves morphological disorders and a stop in reproduction (Alzieu, 2000; Santos, *et al.*, 2002). It also has deleterious effects on fish immune system, thereby increasing fish susceptibility to pathogen infections (Nakayama, *et al.*, 2009).

Despite the ban declared by the International Convention on the Control of Harmful Antifouling Systems on Ships, 2001 (AFS Convention), TBT will remain a concern for several years because this antifouling biocide accumulates in sediments (Bray, 2006; Langston, *et al.*, 2009), which might be re-suspended at one time or another. TBT has a decades long decomposition time, in particular in temperate and cold climates and when in the sediment. High concentrations of TBT have been found worldwide in the vicinity of ports, even in remote areas. For example, in mussels collected near harbors of northern Norway (Kannan & Tanabe, 2009). The 'imposex' phenomenon – i.e. male characteristics appearing in female individuals – has been observed on mussels and snails from the coasts of e.g. Norway, Iceland, the Faroe Islands and Svalbard (OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic, 2000). TBT contamination has also been recorded in Antarctic marine sediments (Negri, *et al.*, 2004). Imposex, along with high levels of TBT and metabolites, has further been demonstrated in densely shipped seas, and the imposex incidence was correlated with the number of ships passing in the vicinity (e.g. ten Hallers-Tjabbes, *et al.*, 1994; ten Hallers-Tjabbes *et al.*, 2003).

The biocidal antifouling agents used since 2008 may also impose problems on the environment and on non-target species. Indeed, even though Cu, which is the biocide of choice for present-day antifouling paints, is an essential micronutrient used in enzymes involved in several metabolic processes, the metal may negatively affect organisms at concentrations higher than physiologically necessary. It has been found that Cu from antifouling paint is an important anthropogenic source of Cu to the aquatic environment and that it in marine environments often exceeds water quality criteria (e.g. Srinivasan, *et al.*, 2007). Apart from the toxicity of copper, synergistic effects when combined with booster biocides may be a problem (e.g. Bao, *et al.*, 2013).

6 INTERNATIONAL BIOFOULING MANAGEMENT MEASURES

International measures

In 2011, the Marine Environment Protection Committee (MEPC) of the IMO adopted the voluntary 'Guidelines for the control and management of ships' biofouling to minimize the transfer of invasive aquatic species' (IMO, 2011). A separate guidance document, based on these Guidelines, provides advice relevant to owners and/or operators of recreational craft less than 24 metres in length, using terminology appropriate for that sector. These guidelines represent the first international action addressing ships' biofouling. The intended goal is to reduce the accumulation of micro- and macro-organisms on the outside of ships by choosing the **appropriate coating**, by conducting **in-water inspections and cleaning**, as well as **proper removal during drydocking**. Ships are required to follow a biofouling management plan and to keep a record book. The guidelines also provide recommendations for the management of **biofouling waste** in land-based facilities. The guidelines will be non-mandatory and as such are not legally enforceable at a global level. However, it is possible that some countries will implement the provisions in the guidance (or parts of them) into national law to protect their waters from invasive marine species from biofouling on ships (IMO, 2011).

In February 2013, the BLG Sub-Committee drafted a guidance document for evaluating the effectiveness of the 2011 Guidelines for the control and management of ships' biofouling. The objective of this initiative is to provide a thorough mechanism for ensuring that the 2011 Biofouling Guidelines are being implemented.

7 EXAMPLES OF SPECIES INTRODUCED INTO THE NORTH SEA THROUGH SHIPS' BIOFOULING

Didemnum vexillum

Didemnum vexillum is a bottom dwelling tunicate native to the waters around Japan. It has however been reported as an invasive species in a number of places in Europe and North America. This species can attach and build over most substrates and will even grow over other organisms when it needs more room to expand (e.g. Bullard *et al.*, 2007; Auken and Oviatt, 2008; Gittenberger, 2007; Valentine *et al.*, 2007; Dijkstra and Harris, 2009). This trait makes it a competitor for resources (e.g. suitable attachment substrates, food, etc.). Its appearance has given rise to its common name marine vomit. Interestingly, this species is not only confined to disturbed and polluted areas, but is also common in the more clean waters where its invasions potentially might have implications on industries in "cleaner" waters, such as fisheries and aquaculture (Bullard *et al.* 2007; Valentine *et al.* 2007), and may further impact natural ecosystems by altering the local habitat (Bullard *et al.* 2007; Valentine *et al.* 2007).



Didemnum vexillum

U.S. Geological Survey/photo by Dann Blackwood

Dasya baillouviana

Dasya baillouviana is a red macroalga. It is a bottom-dweller that usually grows to a length of 50-75 cm. Reproduction can be either asexual or sexual, the latter being more common. Its reproductive cells do not have flagella and it is considered passive because it is not capable of propelling itself in the water and is carried by oceanic currents. *D. baillouviana* can also grow on the shells of bivalves, such as oysters (Haydar & Wolff, 2011) and may therefore compete with other species of algae for space to grow on solid materials. Fertile plants of *D. baillouviana* can develop in just six weeks. *D. baillouviana* is native of the Mediterranean Sea – in particular, along the coasts of Corsica (Coppejans, 1979), the Black Sea – Russian shelf area (Anonymous, 2008) and the Atlantic coast of the United States (Hay & Sutherland, 1988). *D. baillouviana* appeared for the first time in the North Sea at the beginning of the 1950s, in Dutch coastal waters (Stegenga & Prud'homme Van Reine, 1999; Wolff, 2005). Its presence in the Skagerrak was reported along the Swedish west coast in 1953, in Denmark in 1961, and along the south coast of Norway in 1966 (Hopkins, 2002). The species was present in Danish waters in the 1990s (Nielsen, 2005). In 1999, very large specimens of the species were observed in the Kattegat, growing in areas where warm water was discharged by a nuclear power plant (e.g. ICES, 2000; Nyberg, 2007). *D. baillouviana* was spotted in Germany in 2002 (ICES, 2006). It is not sure whether the species was transported from the Western Atlantic through shipping (Hopkins, 2001; ICES, 2006; Gollasch, *et al.*, 2009) or aquaculture activities (Reise, *et al.*, 1999; Wolff, 2005).



Dasya baillouviana growing on the shells of mussels.

Photo: Annelie Lindgren, Dep. of Marine Ecology, Gothenburg University.

Elminius modestus

Elminius modestus is a marine crustacean, a benthic suspension feeder, of the subclass Cirripedia, whose diameter usually measures between 5 and 10 mm. The outside of the shell is flat, and of a white to grey - sometimes darker - colour (OECD, 1963). *E. modestus* can survive up to ten days outside water, tolerates low or variable salinity levels, turbid conditions, low temperatures and even pollution (Crisp, 1958; JNCC, 2012). This barnacle has a high fecundity and a physiological optimum above 20°C (Witte, et al., 2010). *E. modestus* can be found in large numbers on piers, pilings, ships, buoys, stones and even seaweeds in sheltered areas and estuaries. It is less abundant in areas influenced by ocean currents (OECD, 1963). This crustacean rapidly outnumbers native species, such as *Balanus balanoides*, and competes for space with them (Crisp, 1958). *E. modestus* is native of the southern Pacific, particularly of Australia and New Zealand (OECD, 1963; Reise, et al., 1999; Hopkins, 2001; Kerckhof, et al., 2007). It was transported from the Southern Pacific to the United Kingdom on ships' hulls (OECD, 1963; Gollasch, et al., 2009) and possibly in ballast waters during larval stage (Hopkins, 2001; JNCC, 2012). It was first recorded in 1945 in Chichester Harbour, United Kingdom (Crisp, 1958; Reise, et al., 1999). *E. modestus* then spread to France, Belgium, the Netherlands, Germany and Denmark, by both marginal and remote dispersal - i.e. larval drift, favoured by an eastward current, and transfer by ships (Crisp, 1958; Drévès, 2001; Wolf, 2005; Kerckhof, et al., 2007). In addition, *E. modestus* was unintentionally introduced into French bivalve farming areas through the importation of cultivated bivalve molluscs (Pigeot et al., 2001). Although it remained scarce during many years, it developed quickly and outnumbered native barnacles from 2007 onwards. Modelling studies on the effects of climate change and rising temperatures predict that *Elminius modestus* is likely to extend its range in the years to come (Reid et al., 2009).



Elminius modestus, here on wood.

Photo: Christian Buschbaum.

Ficopomatus enigmaticus

Ficopomatus enigmaticus is an invertebrate of the class Polychaeta. The tubeworm grows between 1.5 and 2 cm per month and produces a calcareous tube that serves as a protection (Camus, *et al.*, 2000). Tubes bind together and form reefs which can reach up to several decimeters in thickness (Camus, *et al.*, 2000), and up to 13 kg in weight in three months (ICES, 2009). This species is highly tolerant to variations in its environmental conditions. Although the invertebrate is generally found in brackish waters (Wolff, 2005; Gollasch, *et al.*, 2009), where it has an optimal growth, it can nevertheless tolerate a salinity ranging from 5 to 55 PSU. *F. enigmaticus* can withstand temperatures between 0°C and 35°C and tolerate a pH ranging from 4 to 9. The reproduction and growth of *F. enigmaticus* seem to depend on nutrient abundance, low salinity and weak currents. Indeed, the mechanical action of waves may hamper its development (Schwindt, *et al.*, 2004a). *F. enigmaticus* is considered to be an 'ecosystem engineer' inasmuch as it can have physical impacts on natural systems (Schwindt, *et al.*, 2004b). The species builds reefs whose shape may depend on water depth, water flow direction and the nature of the surface upon which the tubes are attached. When these reefs combine to form one single circular structure, the whole can measure up to 7 m in diameter and 0.5 m in height. It gives shelter to thousands of individual *F. enigmaticus* but also to other species (Schwindt, *et al.*, 2004b). Reef expansion may result in geomorphologic and hydrodynamic alterations in the receiving environment (Schwindt, *et al.*, 2004a). In addition to environmental impacts, the development of *F. enigmaticus* can result in pipe clogging, as well as fouling of port infrastructure and ship hulls (Camus, *et al.*, 2000 ; ICES, 2000, 2006, 2009). *F. enigmaticus* is native of the Indian Ocean and Southern Pacific (Reise, *et al.*, 1999 ; Hopkins, 2001; Wolff, 2005 ; ICES, 2009). *F. enigmaticus* was observed in France, in 1921 (Camus, *et al.* 2000 ; ICES, 2000, 2009), soon after in United Kingdom in 1922 (Gollasch, *et al.*, 2009) and in the Netherlands in 1968 (Wolff, 2005). The polychaete was also reported in Denmark (ICES, 2009). It is considered established in United Kingdom, Germany, the Netherlands (Wolff, 2005) and Belgium (Gollasch, *et al.*, 2009). It is unclear whether *F. enigmaticus* was introduced by shipping (Reise, *et al.*, 1999 ; Hopkins, 2001 ; Wolff, 2005) or aquaculture activities (Gollasch, *et al.*, 2009).



Ficopomatus enigmaticus

Photo credits: Museo di Storia Naturale di Venezia.

Laminaria ochotensis

Laminaria ochotensis is an edible brown alga (kelp) of the kingdom Phaeophyta which grows at a water depth of about 3 to 10 m (Miyabe, 1902). *L. ochotensis* lives naturally along the coasts of Japan (Wallentinus, 2002; Selivanova, *et al.*, 2007). The species has been observed in Germany and its introduction has been attributed to hull fouling (Selivanova, *et al.*, 2007; Gollasch, *et al.*, 2009).

8 CONCLUSION

The issue of alien species translocation in biofouling is being dealt with in IMO Guidelines. The existing IMO Guidelines however need to be updated regarding for example niche areas on ships, such as sea chests, as these enclosed small spaces with lack of effective antifouling paints and elevated temperatures provide suitable conditions for a variety of species and larger adult marine organisms that might not survive on a hull surface or in ballast water.

Policy makers and others involved in developing policies for biofouling have spent ample thoughts on a feasible form for regulating biofouling. The question is would it best fit under another Convention, notably the Anti-Fouling Convention (2001) or the Ballast Water Management Convention (2004), or be a self-standing instrument?

9 References

- Ahimou, F., Semmens, M.J., Haugstad, G. & Novak, P.J. (2007). Effect of protein, polysaccharide, and oxygen concentration profiles on biofilm cohesiveness. *Applied and Environmental Microbiology*. Vol. 73. No. 9.
- Allison, D.G., Maira-Litran, T. & Gilbert, P. (2000). Antimicrobial resistance of biofilms. In L.V. Evans (Ed.), *Biofilms : recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.
- Alzieu, C. (2000, April 14). *Environmental impact of TBT : the French experience*. The Science of the Total Environment 258 (2000). Elsevier.
- Anonymous. (2008). Commission on the Protection of the Black Sea against Pollution. In Oguz, T.(Ed.) *State of the Environment of the Black Sea (2001-2006/7)*.
- Arai, T. (2009). Preface. In T. Arai, M. Ohji, H. Harino & W.J. Langston (Eds.), *Ecotoxicology of antifouling biocides*. Springer. Tokyo.
- Auker, L.A. & Oviatt, C.A. (2008) Factors influencing the recruitment and abundance of *Didemnum* in Narragansett Bay, Rhode Island., *ICES Journal of Marine Science* 65: 765-769
- Bao, V.W.W., Leung, K.M.Y., Lui, G.C.S. & Lam, M.H.W. (2013). Acute and chronic toxicities of Irgarol alone and in combination with copper to the marine copepod *Tigriopus japonicas*. *Chemosphere*. Vol. 90 3:1140-1148.
- Bray, S. (2006). *Tributyltin pollution on a global scale : an overview of relevant and recent research : impacts and issues*. Edited by W. J. Langston, Marine Biological Association, Plymouth, United Kingdom. http://assets.wwf.no/downloads/tbt_global_review_wwf_uk_oct_2006.pdf
- Bullard, S. G. and 15 authors. (2007) The colonial ascidian *Didemnum* sp. A: Current distribution, basic biology and potential threat to marine communities of the northeast and west coasts of North America., *Journal of Experimental Marine Biology and Ecology* 342: 99-108
- Callow, M.E. (2000). Algal biofilms. In L.V. Evans (Ed.), *Biofilms: recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.
- Camus, P., Compère, C., Blanchet, A., Dimeet, J., Hamon, D., Lacotte, N., Peleau, M. & Lassalle, E. (2000). *Ficopomatus enigmaticus* : écologie, répartition en France et Bretagne, nuisances et moyens de lutte sur le site atelier du port de Vannes (résumé). <http://envlit.ifremer.fr/content/download/27417/222402/version/1/file/triniteficoresume.pdf>
- Callow, M.E. & Callow, J.A. (2002). Marine biofouling : a sticky problem. *Biologist*. Vol. 49. No. 1.
- Callow, J.A. & Callow, M.E. (2011). Trends in the development of environmentally friendly fouling-resistant marine coatings. *Nature Communications*. 2:244.
- Camus, P., Compère C., Blanchet A., Dimeet J., Hamon D., Lacotte N., et al. (2000). *Ficopomatus enigmaticus*: écologie, répartition en France et Bretagne, nuisances et moyens de lutte sur le site atelier du port de Vannes (résumé).
- Carlton, J.T. (1996). Marine bioinvasions : the alteration of marine ecosystems by non-indigenous species. *Oceanography*. Vol. 9. No. 1. http://www.tos.org/oceanography/issues/issue_archive/issue_pdfs/9_1/9.1_carlton.pdf
- Chambers, L.D., Stokes, K.R., Walsh, F.C. & Wood, R.J.K. (2006). Modern approaches to marine antifouling coatings. *Surface & Coatings Technology*. Vol. 201.
- Champ, M.A. (2001, November 8). *The status of the treaty to ban TBT in marine antifouling paints and alternatives*. Proceedings of the 24th UJNR (US/Japan) Marine Facilities Panel Meeting in Hawaii. National Maritime Research Institute of Japan. http://www.nmri.go.jp/main/cooperation/ujnr/24ujnr_paper_us/Ocean_Engineering_and_Resource/OERD_Champ.pdf
- Coppejans, E. (1979). Végétation marine de la Corse (Méditerranée). Documents pour la flore des algues. *Botanica Marina* XX11(4).
- Coutts ADM, Moore KM, Hewitt CL (2003). Ships' sea-chests: an overlooked transfer mechanism for non-indigenous marine species? *Marine Pollution Bulletin* 46: 1504–1515. [http://dx.doi.org/10.1016/S0025-326X\(03\)00292-3](http://dx.doi.org/10.1016/S0025-326X(03)00292-3)
- Coutts ADM, Taylor MD (2004). A preliminary investigation of biosecurity risks associated with biofouling on merchant vessels in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 38: 215–229. <http://dx.doi.org/10.1080/00288330.2004.9517232>
- Coutts ADM, Dodgshun TJ (2007). The nature and extent of organisms in vessel sea-chests: A protected mechanism for marine bioinvasions. *Marine Pollution Bulletin* 54: 875–886, <http://dx.doi.org/10.1016/j.marpolbul.2007.03.011>
- Crisp, D. J. (1958). The spread of *Elminius modestus* Darwin in north-west Europe. *Journal of the Marine Biological Association of the United Kingdom*. 37(2).
- Dijkstra, J.A.; Harris, L.G. (2009) Maintenance of diversity altered by a shift in dominant species: implications for species coexistence, *Marine Ecology Progress Series* 387: 71-80.

- Donlan, R.M. (2000). Biofilm control in industrial water systems : approaching an old problem in new ways. In L.V. Evans (Ed.), *Biofilms: recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.
- Drake, J.M. & Lodge, D.M. (2007, May). Hull fouling is a risk factor for intercontinental species exchange in aquatic ecosystems. *Aquatic Invasions*. Vol. 2. No. 2.
- Drévès, L. (2001). Effets climatiques sur les écosystèmes marins : exemple du recrutement des crustacés cirripèdes sur la côte ouest du Cotentin. *Hydroécologie Appliquée*. 13.
- Flemming, H.-C., Wingender, J., Griebe, T. & Mayer, C. (2000). Physico-chemical properties of biofilms. In L.V. Evans (Ed.), *Biofilms : recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.
- Flemming, H.C. (2002). Biofouling in water systems : cases, causes and countermeasures. *Applied Microbiology and Biotechnology*. Vol. 59. No. 6.
- Frey, M.A., Simard, N., Robichaud, D.D., Martin, J.L. & Therriault, T.W. (2014). Fouling around: vessel sea-chests as vector for the introduction and spread of aquatic invasive species. *Management of biological invasions*, Vol 5. 1:21-30.
- Gittenberger, A. (2007) Recent population expansions of non-native ascidians in the Netherlands., *Journal of Experimental Marine Biology and Ecology* 342: 122-126.
- Gollasch, S. (2002). The importance of ship hull fouling as a vector of species introductions into the North Sea. *Biofouling*. Vol. 18. No. 2.
- Gollasch, S. (2006, October 30). *Odontella sinensis*. Delivering Alien Invasive Species Inventories for Europe. http://www.europe-aliens.org/pdf/Odontella_sinensis.pdf
- Gollasch, S., Haydar, D., Minchin, D., Wolff, W.J. & Reise, K. (2009). Introduced Aquatic Species of the North Sea Coasts and Adjacent Brackish Waters. In G. Rilov & J.A. Crooks (Eds.) *Biological Invasions in Marine Ecosystems*. Ecological Studies 204. Springer-Verlag Berlin. <http://www.rug.nl/biologie/onderzoek/overonderzoek/onderzoekgroepen/marbee/publications/2009/GollaschEcolStud09.pdf>
- Hay, M.E., & Sutherland J.P. (1988). The ecology of rubble structures of the South Atlantic bight: a community profile. Performed for U.S. Department of the Interior, Fish and Wildlife Service, National Wetlands Research Center.
- Haydar, D., & Wolff W.J. (2011). Predicting invasion patterns in coastal ecosystems: relationship between vector strength and vector tempo. *Marine Ecology Progress Series*. 431.
- Hellio, C., Tsoukatou, M., Maréchal, J-P., Beaupoil, C., Clare, A.S., Vagias, C. & Roussis, V. (2005). Inhibitory effects of Mediterranean sponge extracts and metabolites on larval settlement of the barnacle *Balanus amphitrite*. *Marine Biotechnology*, Vol. 7:297-305.
- Holmström, C. & Kjelleberg, S. (2000). Bacterial interactions with marine fouling organisms. In L.V. Evans (Ed.), *Biofilms : recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.
- Hopkins, G.A. & Forrest, B.M. (2008). Management options for vessel hull fouling: an overview of risks posed by in-water cleaning. *ICES J. Mar. Sci.* (2008) 65 (5):811-815.
- Hopkins, C.C.E. (2001, February). *Actual and potential effects of introduced marine organisms in Norwegian waters, including Svalbard*. Research report 2001-1. Directorate for Nature Management.
- Hopkins, C.C.E. (2002). Introduced marine organisms in Norwegian waters, including Svalbard. In E. Leppäkoski, S. Gollasch & S. Olenin (Eds.), *Invasive aquatic species of Europe : distribution, impacts and management*. Kluwer Academic Publishers. Dordrecht. The Netherlands.
- International Council for the Exploration of the Sea (ICES). (2000). *Report of the working group on introductions and transfers of marine organisms*. Parnu, Estonia, 27-29 March, 2000. <http://www.ices.dk/reports/acme/2000/wgitmo00.pdf>
- International Council for the Exploration of the Sea (ICES). (2006). *Working Group on Introductions and Transfers of Marine Organisms (WGITMO)*. 16-17 March 2006, Oostende, Belgium. <http://www.dirnat.no/multimedia/1517/WGITMO.pdf&contentdisposition=attachment>
- International Council for the Exploration of the Sea (ICES). (2009). *Report of the Working Group on Introduction and Transfers of Marine Organisms (WGITMO)*. 11 - 13 March 2009, Washington D.C., USA. <http://www.ices.dk/reports/ACOM/2009/WGITMO/WGITMO2009.pdf>
- International Maritime Organization. (2010, July 23). *Reduction of GHG emissions from ships : marginal abatement costs and cost-effectiveness of energy-efficiency measures*. Submitted by the Institute of Marine Engineering, Science and Technology (MEPC 61/INF.18). London. <http://docs.imo.org/>
- International Maritime Organization. (2011, July 26). *Report of the Marine Environment Protection Committee on its sixty-second session* (MEPC 62/24/Add.1). London. <http://docs.imo.org/>
- Jain, A. & Bhosle, N. B. (2009) Biochemical composition of the marine conditioning film: implications for bacterial adhesion. *Biofouling* 25, 13-19.
- Joint Nature Conservation Committee (JNCC). (2012). *Elminius modestus*. Retrieved June 18, 2012, from <http://jncc.defra.gov.uk/page-1704>

- Kannan, K. & Tanabe, S. (2009). Global contamination by organotin compounds. In T. Arai, H. Harino, M. Ohji & W.J. Langston (Eds.), *Ecotoxicology of antifouling biocides*. Springer. Tokyo.
- Kerckhof, F., Haelters J., & Gollasch S. (2007). Alien species in the marine and brackish ecosystem: the situation in Belgian waters. *Aquatic Invasions*. 2(3): 243 - 257.
- Kim, T.W. & Micheli, F. (2013) Decreased solar radiation and increased temperature combine to facilitate fouling by marine non-indigenous species. *Biofouling*. Vol 29. No. 5, 501-512.
- Kooistra, W.H.C.F., De Boer, M.K., Vrieling, E.G., Connell, L.B. & Gieskes, W.W.C. (2001). Variation along ITS markers across strains of *Fibrocapsa japonica* (Raphidophyceae) suggests hybridization events and recent range expansion. *Journal of Sea Research*. Vol. 46. Elsevier.
- Krøber, A.C., Carstens, K., Peters, S., Tilly, K. & Wiltshire, K.H. (2012, September). The diatom *Mediopyxis helysia* (Kühn, Hargreaves & Halliger 2006) at Helgoland Roads : a success story ? *Helgoland Marine Research*. Vol. 66. Issue 3.
- Kubaneck, J., Whalen, K.E., Engel, S., Kelly, S.R., Henkel, T.P., Fenical, W. & Pawfik, J.R. (2002). Multiple defensive roles for triterpene glycosides from two Caribbean sponges. *Oecologia* 1: 125-136.
- Langston, W.J., Harino, H. & Pope, N.D. (2009). Behaviour of organotins in the coastal environment. In T. Arai, H. Harino, M. Ohji & W.J. Langston (Eds.), *Ecotoxicology of antifouling biocides*. Springer. Tokyo.
- Lewandowski, Z. (2000). Structure and function of biofilms. In L.V. Evans (Ed.), *Biofilms : recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.
- Meesters, K.P.H., Van Groenestijn, J.W. & Gerritse, J. (2003). Biofouling reduction in recirculating cooling systems through biofiltration of process water. *Water Research*. Vol. 37. No. 3.
- Mineur, F., Johnson, M.P., Maggs, C.A. & Stegenga, H. (2007). *Hull-fouling on commercial ships as a vector of macro algal introduction*. Marine Biology. Volume 151. Springer-Verlag. <http://fn3dn7qc5p.scholar.serialssolutions.com/>
- Ministry of Agriculture and Forestry Biosecurity New Zealand. (2010a, March). *Vessel biofouling as a vector for the introduction of non-indigenous marine species to New Zealand : fishing vessels*. MAF Biosecurity New Zealand Technical Paper No: 2010/11.
- Ministry of Agriculture and Forestry Biosecurity New Zealand. (2010b, July). *Vessel biofouling as a vector for the introduction of non-indigenous marine species to New Zealand : slow-moving barges and oil platforms*. Biosecurity New Zealand Project ZBS2005-02. MAF Biosecurity New Zealand Technical Paper No: 2010/12.
- Miyabe, K. (1902). On the Laminariaceae of Hokkaido. The Journal of the Sapporo Agricultural College. Hokkaido University Collection of Scholarly and Academic Papers.
- Nakayama, A., Segner, H. & Kawai, S. (2009). Immunotoxic effects of organotin compounds in teleost fish. In T. Arai, H. Harino, M. Ohji & W.J. Langston (Eds.), *Ecotoxicology of antifouling biocides*. Springer. Tokyo.
- Negri, A.P., Hales, L.T., Battershill, C., Wolff, C. & Webster, N.S. (2004). TBT contamination identified in Antarctic marine sediments. *Marine Pollution Bulletin*. Vol. 48. No. 11-12.
- Organisation for economic co-operation and development. (1963). *Catalogue of main marine fouling organisms found on ships coming into European waters*. Volume 1. Barnacles.
- Nielsen, R. (2005). Danish seaweeds. Botanical Museum, Copenhagen, Denmark.
- Nyberg, C. D. (2007). Introduced marine macroalgae and habitat modifiers: their ecological role and significant attributes. Department of Marine Ecology. Göteborg University. Sweden. Ph.D.thesis.
- OECD (1963). *Catalogue of main marine fouling organisms found on ships coming into European waters*. Barnacles. 1.
- Orr, R.J.S., Stüken, A., Rundberget, T., Eikrem W., & Jakobsen K. J. (2011). Improved phylogenetic resolution of toxic and non-toxic Alexandrium strains using a concatenated rDNA approach. *Harmful Algae*. 10.
- OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic. (2000). *Quality Status Report 2000*. Arctic Waters. Chapter 4 Chemistry. <http://www.ospar.org/>
- Patin, S. (1999). Environmental impact of the offshore oil and gas industry. EcoMonitor Publishing. New York.
- Pigeot, J., Miramand P., Garcia-Meunier P., Guyot T., & Séguignes M. (2000). Présence d'un nouveau prédateur de l'huître creuse, *Ocenebrellus inornatus* (Récluz, 1851), dans le bassin conchylicole de Marennes-Oléron. *Comptes rendus (sciences de la vie) de l'Académie des sciences de Paris*. 323.
- Quiniou, C. & Compère, C. (2009). *La chimie et la mer : ensemble au service de l'homme*. EDP Sciences. Les Ulis (France).
- Railkin, A.I. (2004). *Marine biofouling : colonization processes and defenses*. CRC Press.
- Readman, J.W., Van Hattum, B., Barcelo, D., Albanis, T.A., Riemann, B., Blanck, H., Gustavson, K., Tronczynski, J. & Jacobson, A. (2002, 27 June). *Assessment of antifouling agents in coastal environments (ACE): Final Scientific and Technical Report*. MAS3-CT98-0178. <http://www.pml-ace.org.uk/>
- Reid, P. C., Cook, E. J., Edwards, M., McQuatters-Gollop, A., Minchin, D., & McCollin, T. (2009). Marine non-native species. Marine Climate Change Impacts Partnership Ecosystem Linkages Report Card 2009 Science Review.

- Reise, K., Gollasch, S. & Wolff, W.J. (1999). *Introduced marine species of the North Sea coasts*. Helgoländer Meeresuntersuchungen. Helgoländer Meeresunters 52. Biologische Anstalt Helgoland, Hamburg. http://gollaschconsulting.de/download/1999_invaders_north_sea_kr_sg_ww.pdf
- Russell, A.D. (2003, December). Biocide use and antibiotic resistance: the relevance of laboratory findings to clinical and environmental situations. *The Lancet Infectious Diseases*. Volume 3. Issue 12.
- Santos, M.M., Ten Hallers-Tjabbes, C.C., Santos, A.M. & Vieira, N. (2002). Imposex in *Nucella lapillus*, a bioindicator for TBT contamination : re-survey along the Portuguese coast to monitor the effectiveness of EU regulation. *Journal of Sea Research*. Vol. 48. No. 3.
- Schwindt, E., Iribarne O. O., & Isla F. I. (2004a). Physical effects of an invading reef-building polychaete on an Argentinean estuarine environment. *Estuarine, Coastal and Shelf Science*. 59.
- Schwindt, E., De Francesco, C. G., & Iribarne, O. O. (2004b). Individual and reef growth of the invasive reef-building polychaete *Ficopomatus enigmaticus* in a south-western Atlantic coastal lagoon. *Journal of the Marine Biological Association of the United Kingdom*.
- Sera, Y., Adachi, K., Shizuri, Y. (1999) A new epidioxysterol as an antifouling substance from a Palauan marine sponge, *Lendenfeldia chondrodes*. *J Nat Prod* 62: 152–154.
- Schultz, M.P., Bendick, J.A., Holm, E.R. & Hertel, W.M. (2011). Economic impact of biofouling on a naval surface ship. *Biofouling*. Vol. 27. No. 1.
- Selivanova, O. N., Zhigadlova G. G., & Hansen G. I. (2007). Revision of the Systematics of Algae in the Order Laminariales (Phaeophyta) from the Far-Eastern Seas of Russia on the Basis of Molecular–Phylogenetic Data. *Russian Journal of Marine Biology*. 33(5).
- Srinivasan, M. & Swain, G.W. (2007). Managing the use of copper-based antifouling paints. *Environmental Management*. Vol. 39, 3:423–441.
- Sorte, C.J.B., Williams, S.L. & Zerebecki, R.A. (2010). Ocean warming increases threat of invasive species in a marine fouling community. *Ecology*. Vol. 91. No. 8.
- Stegenga, H. & Prud'homme Van Reine, W.F. (1999). *Changes in the seaweed flora of the Netherlands*. Research Institute Rijksherbarium / Hortus Botanicus. Leiden. The Netherlands. <http://www.vliz.be>
- Swain, G., Herpe, S., Ralston, E. & Tribou, M. (2006). Short-term testing of antifouling surfaces: the importance of colour. *Biofouling*. Vol. 22. No. 5–6.
- Sylvester F, MacIsaac HJ (2010). Is vessel hull fouling an invasion threat to the Great Lakes? *Diversity and Distributions* 16: 132–143. <http://dx.doi.org/10.1111/j.1472-4642.2009.00622.x>
- Tang, Y.Z., Kong L., & Holmes M. J. (2007). Dinoflagellate *Alexandrium leei* (Dinophyceae) from Singapore coastal waters produces a water-soluble ichthyotoxin. *Marine Biology*. 150.
- ten Hallers-Tjabbes, C.C., Kemp, J.F., Boon, J.P. (1994). Imposex in whelks (*Buccinum undatum*) from the open North Sea: relation to shipping traffic intensities. *Mar Pollut Bull*. 28:311–3.
- ten Hallers-Tjabbes, C.C., Wegener, J.-W., van Hattum, B., Kemp, J.F., ten Hallers, E., Reitsema, T. & Boon, J.P. (2003). Imposex and organotin concentrations in *Buccinum undatum* and *Neptunea antiqua* from the North Sea: relationship to shipping density and hydrographical conditions. *Marine Environmental Research*. 55:203–233.
- Valentine, Page C.; Collie, Jeremy S.; Reid, Robert N.; Asch, Rebecca G.; Guida, Vince G.; Blackwood, Dann S. (2007) The occurrence of the colonial ascidian *Didemnum* sp. on Georges Bank gravel habitat — Ecological observations and potential effects on groundfish and scallop fisheries., *Journal of Experimental Marine Biology and Ecology* 342: 179–181
- Verveen, A., Vincx, M. & Degraer, S. (2007). The effect of temperature and salinity on the survival of *Mytilopsis leucophaeata* larvae (Mollusca, Bivalvia): The search for environmental limits. *Journal of Experimental Marine Biology and Ecology*. 348, 111–120.
- Wallentinus, I. (2002). Marine algae and vascular plants in European aquatic environments. In E. Leppäkoski, S. Gollasch & S. Olenin (Eds.), *Invasive aquatic species of Europe : distribution, impacts and management*. Kluwer Academic Publishers. Dordrecht. The Netherlands.
- Witte, S., Buschbaum, C., Beusekom J. E. E., & Reise, K. (2010). Does climatic warming explain why an introduced barnacle finally takes over after a lag of more than 50 years? *Biological Invasions*. 12(10):3579– 3589.
- Wolff, W.J. (2005). Non-indigenous marine and estuarine species in the Netherlands. *Zoologische Mededelingen*. 79(1). <http://www.repository.naturalis.nl/document/41874>
- Zinn, M., Zimmerman, R.C. & White, D.C. (2000). Environmentally acceptable control of microbial biofilms. In L.V. Evans (Ed.), *Biofilms: recent advances in their study and control*. Harwood Academic Publishers. Amsterdam.